

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/12, C12P 21/02, C07K 13/00, C12N 5/10, A61K 37/02	A1	(11) International Publication Number: WO 94/26892 (43) International Publication Date: 24 November 1994 (24.11.94)
(21) International Application Number: PCT/US94/05288 (22) International Filing Date: 12 May 1994 (12.05.94) (30) Priority Data: 08/061,464 12 May 1993 (12.05.93) US (71) Applicant: GENETICS INSTITUTE, INC. [US/US]; Legal Affairs, 87 Cambridge Park Drive, Cambridge, MA 02140 (US). (72) Inventors: CELESTE, Anthony, J.; 86 Packard Street, Hudson, MA 01749 (US). WOZNEY, John, M.; 59 Old Bolton Road, Hudson, MA 01749 (US). (74) Agent: LAZAR, Steven, R.; Genetics Institute, Inc., Legal Affairs, 87 Cambridge Park Drive, Cambridge, MA 02170 (US).		(81) Designated States: AU, BR, CA, FI, JP, KP, KR, NL, NO, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: BMP-11 COMPOSITIONS (57) Abstract Purified BMP-11 proteins and processes for producing them are disclosed. Recombinant DNA molecules encoding the BMP-11 proteins are also disclosed. The proteins may be useful in regulating follicle stimulating hormone, such as for contraception. In addition, the proteins may be useful for the induction of bone, cartilage and/or other connective tissue.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TITLE OF THE INVENTION

BMP-11 COMPOSITIONS

BACKGROUND OF THE INVENTION

The present invention relates to a novel family of purified proteins designated BMP-11, DNA molecules encoding them, and processes for obtaining them. The inventors have previously designated the BMP-11 proteins as Activin WC. The BMP-11 proteins may be useful to induce bone and/or cartilage formation and in wound healing and tissue repair, or for augmenting the activity of other bone morphogenetic proteins. The BMP-11 proteins may also be useful to regulate the production of follicle stimulating hormone, for contraception, to promote neuronal cell survival, to stimulate hematopoiesis, and to suppress the development of gonadal tumors.

United States Patent 4,798,885 disclosed DNA encoding the prepro inhibin α and β chains. United States Patent 5,071,834 discloses pharmaceutical compositions of activin with two beta₂ chains formulated in a pharmaceutically acceptable carrier.

United States Patent 5,102,807 discloses a purified inhibin protein which suppresses production of FSH without suppressing production of luteinizing hormone.

SUMMARY OF THE INVENTION

BMP-11 protein is a member of the TGF- β superfamily of proteins. The TGF- β superfamily includes the family of proteins known as bone morphogenetic proteins (BMPs), as well as a group of proteins that are termed inhibin- β . As discussed further herein, when dimerized with another BMP-11 (homodimer), BMP-11 protein is expected to demonstrate BMP-11 activity, as further described herein, as may be measured in accordance with the assays described in the examples herein. When dimerized as a heterodimer with inhibin- α proteins or with other inhibin- β proteins, the inhibin- β /BMP-11 heterodimer is expected to demonstrate effects on the production of follicle stimulating hormone (FSH), as described further herein. It is further expected that, in homodimeric form or in heterodimeric form with another member of the bone morphogenetic protein family, BMP-11

will exhibit BMP activity, i.e., the ability to induce the formation of bone, cartilage and/or other connective tissue. Thus, depending upon the environment of BMP-11, it may form dimers which will demonstrate either activin or inhibin activity, or bone, cartilage and/or other connective tissue-inducing activity. Accordingly, BMP-11 activity is defined as the ability to regulate the production of FSH in the assay described at Example 8 herein, or the ability to induce the formation of bone, cartilage and/or other connective tissue in the assays described at Examples 5 to 7 herein.

Proteins termed inhibins and activins are produced in the gonad and exist naturally in follicular fluid. These proteins act at the level of the anterior pituitary gland to inhibit (inhibins) or stimulate (activins) the release of follicle-stimulating hormone (FSH) [for reviews see, e.g., Ying, S.-Y., *Endocr. Rev.*, 9:267-293 (1988) or Ling, N. et al, *Vitamins and Hormones*, 44:1-46 (Academic Press 1988)]. Briefly, dimeric proteins comprised of one chain of inhibin α and one chain of inhibin β (β_A or β_B) are termed inhibins and are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while other dimeric proteins comprised of two chains of inhibin β (β_A or β_B) are termed activins and are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH) [see, e.g., Ling et al., *Nature*, 321:779-782 (1986) or Vale, et al., *Nature*, 321:776-779 (1986) or Mason et al., *Nature*, 318:659-663 (1985) or Forage et al., *Proc. Natl. Acad. Sci. USA*, 83:3091-3095 (1986)].

It is recognized that FSH stimulates the development of ova in mammalian ovaries (Ross et al., in *Textbook of Endocrinology*, ed. Williams, p. 355 (1981) and that excessive stimulation of the ovaries with FSH will lead to multiple ovulations. FSH is also important in testicular function. Thus, BMP-11, in heterodimers with a member of the inhibin α family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility these mammals.

BMP-11, as a homodimer or as a heterodimer with other protein subunits of the inhibin- β group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. BMP-11 may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs. It is further contemplated that BMP-11 may be useful in promoting neuronal cell survival [see, e.g., Schubert et al., *Nature*, 344:868-870 (1990)], modulating hematopoiesis by inducing the differentiation of erythroid cells [see, e.g., Broxmeyer et al, *Proc. Natl. Acad. Sci. USA*, 85:9052-9056 (1988) or Eto et al, *Biochem. Biophys. Res. Comm.*, 142:1095-1103 (1987)], for suppressing the development of gonadal tumors [see, e.g., Matzuk et al., *Nature*, 360:313-319 (1992)] or for augmenting the activity of bone morphogenetic proteins [see, e.g., Ogawa et al., *J. Biol. Chem.*, 267:14233-14237 (1992)].

BMP-11 proteins may be further characterized by their ability to modulate the release of follicle stimulating hormone (FSH) in established *in vitro* bioassays using rat anterior pituitary cells as described [see, e.g., Vale et al, *Endocrinology*, 91:562-572 (1972); Ling et al., *Nature*, 321:779-782 (1986) or Vale et al., *Nature*, 321:776-779 (1986)]. It is contemplated that the BMP-11 protein of the invention, when composed as a homodimer or a heterodimer with other inhibin β chains will exhibit stimulatory effects on the release of follicle stimulating hormone (FSH) from anterior pituitary cells as described [Ling et al., *Nature*, 321:779-782 (1986) or Vale et al., *Nature*, 321:776-779 (1986)]. Additionally, it is contemplated that the BMP-11 protein of the invention, when composed as a heterodimer with the inhibin α chain, will inhibit the release of follicle stimulating hormone (FSH) from anterior pituitary cells as described [see, e.g., Vale et al, *Endocrinology*, 91:562-572 (1972)]. Therefore, depending on the particular composition, it is expected that the BMP-11 protein

of the invention may have contrasting and opposite effects on the release of follicle stimulating hormone (FSH) from the anterior pituitary.

5 Activin A (the homodimeric composition of inhibin β_A) has been shown to have erythropoietic-stimulating activity [see e.g. Eto et al., *Biochem. Biophys. Res. Commun.*, 142:1095-1103 (1987) and Murata et al., *Proc. Natl. Acad. Sci. U.S.A.*, 85:2434-2438 (1988) and Yu et al., *Nature*, 330:765-767 (1987)]. It is contemplated that the BMP-11 protein of the invention has a
10 similar erythropoietic-stimulating activity. This activity of the BMP-11 protein may be further characterized by the ability of the BMP-11 protein to demonstrate erythropoietin activity in the biological assay performed using the human K-562 cell line as described by [Lozzio et al., *Blood*, 45:321-334 (1975) and
15 U.S. Pat. No. 5,071,834].

 The structures of several proteins, designated BMP-1 through BMP-9, have previously been elucidated. The unique inductive activities of these proteins, along with their presence in bone, suggests that they are important regulators of
20 bone repair processes, and may be involved in the normal maintenance of bone tissue. The BMP-11 protein of the present invention is related to the above BMP proteins, and is expected to share BMP activities such as the ability to induce bone, cartilage and/or other connective tissue, such as tendon or
25 ligament, and wound healing activities of the BMPs. In addition, it is expected that the proteins of the invention may act in concert with or perhaps synergistically with other related proteins and growth factors. Further therapeutic methods and compositions of the invention therefore comprise a
30 therapeutic amount of at least one BMP-11 protein of the invention with a therapeutic amount of at least one of the other BMP proteins disclosed in co-owned patents and applications described below. Such combinations may comprise separate molecules of the BMP proteins or heteromolecules comprised of
35 different BMP moieties. Further, BMP-11 proteins may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question.

These agents include various growth factors such as epidermal growth factor (EGF), fibroblast growth factor (FGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- α and TGF- β), and k-fibroblast growth factor (kFGF), parathyroid hormone (PTH), leukemia inhibitory factor (LIF/HILDA/DIA), insulin-like growth factors (IGF-I and IGF-II). Portions of these agents may also be used in compositions of the present invention.

The bovine BMP-11 DNA sequence (SEQ ID NO: 1) and amino acid sequence (SEQ ID: 2) and human BMP-11 DNA sequence (SEQ ID NO:10) and amino acid sequence (SEQ ID NO:11) are set forth in the Sequence Listings herein. Activin proteins are capable of regulating the production of follicle-stimulating hormone (FSH), and thus BMP-11 may be useful as a contraceptive or a fertility inducing therapeutic. In homodimeric form or in heterodimers with proteins of the inhibin- β group, purified BMP-11 protein is expected to demonstrate activin activity, and may be used to stimulate FSH. In addition, it is expected that the purified BMP-11 protein may be useful for the induction of bone, cartilage and/or other connective tissue.

Bovine BMP-11 may be produced by culturing a cell transformed with a DNA sequence comprising nucleotide #375 to nucleotide #704 as shown in SEQ ID NO: 1 and recovering and purifying from the culture medium a protein characterized by the amino acid sequence comprising amino acid # 1 to # 109 as shown in SEQ ID NO: 2 substantially free from other proteinaceous materials with which it is co-produced.

Human BMP-11 is expected to be homologous to bovine BMP-11. The invention, therefore, includes methods for obtaining the DNA sequences encoding human BMP-11, the DNA sequences obtained by these methods, and the human protein encoded by these DNA sequences. This method entails utilizing the bovine BMP-11 nucleotide sequence or portions thereof to design probes to screen libraries for the human gene or fragments thereof using standard techniques. A DNA sequence encoding part of the human BMP-11 protein (SEQ ID NO: 3) and the corresponding amino acid sequence (SEQ ID NO: 4) are set forth in the Sequence Listing.

These sequences may also be used in order to design probes to obtain the complete human BMP-11 gene through standard techniques. Human BMP-11 may be produced by culturing a cell transformed with the BMP-11 DNA sequence and recovering and purifying BMP-11 from the culture medium. The purified expressed protein is substantially free from other proteinaceous materials with which it is co-produced, as well as from other contaminants.

The recovered purified protein is contemplated to demonstrate the ability to regulate the production of FSH. The proteins of the invention may be further characterized by the ability to regulate the production of follicle stimulating hormone (FSH) in established in vitro bioassays using rat anterior pituitary cells. BMP-11 proteins may also be characterized by the ability to induce the formation of bone, cartilage and/or other connective tissue, for example, in the rat bone formation assay described below.

Another aspect of the invention provides pharmaceutical compositions containing a therapeutically effective amount of a BMP-11 protein in a pharmaceutically acceptable vehicle or carrier. BMP-11 compositions of the invention may be useful for the regulation of follicle stimulating hormone, and may be useful in contraception. Compositions of the invention may further include at least one other therapeutically useful agent such as the BMP proteins BMP-1, BMP-2, BMP-3, BMP-4, BMP-5, BMP-6 and BMP-7, disclosed for instance in United States Patents 5,108,922; 5,013,649; 5,116,738; 5,106,748; 5,187,076; and 5,141,905; BMP-8, disclosed in PCT publication WO91/18098; and BMP-9, disclosed in PCT publication WO93/00432, and BMP-10, disclosed in co-pending patent application serial number 08/061,695, filed on May 12, 1993. The BMP-11 compositions may also be useful for a number of uses involving regulation of the production of follicle stimulating hormone, including contraception. These methods, according to the invention, entail administering to a patient needing such treatment, an effective amount of BMP-11.

The compositions of the invention may comprise, in addition

to a BMP-11 protein, other members of the inhibin- β group of proteins or inhibin- α proteins, as well as other therapeutically useful agents including growth factors such as epidermal growth factor (EGF), fibroblast growth factor (FGF), transforming growth factor (TGF- α and TGF- β), and insulin-like growth factor (IGF).

The BMP-11 compositions of the present invention may also be useful for treating a number of bone and/or cartilage defects, periodontal disease and various types of wounds. These methods, according to the invention, entail administering to a patient needing such bone and/or cartilage formation wound healing or tissue repair, an effective amount of a BMP-11 protein. These methods may also entail the administration of a protein of the invention in conjunction with at least one of the novel BMP proteins disclosed in the co-owned patents and applications described above. In addition, these methods may also include the administration of a BMP-11 protein with other growth factors including EGF, FGF, TGF- α , TGF- β , and IGF.

Still a further aspect of the invention are DNA sequences coding for expression of a BMP-11 protein. Such sequences include the sequence of nucleotides in a 5' to 3' direction illustrated in SEQ ID NO: 1 or DNA sequences which hybridize under stringent conditions with the DNA sequence of SEQ ID NO: 1 and encode a protein having BMP-11 activity. Finally, allelic or other variations of the sequences of SEQ ID NO: 1, whether such nucleotide changes result in changes in the peptide sequence or not, are also included in the present invention.

Still a further aspect of the invention are DNA sequences coding for expression of a BMP-11 protein. Such sequences include the sequence of nucleotides in a 5' to 3' direction illustrated in SEQ ID NO: 1 or SEQ ID NO:10, and DNA sequences which, but for the degeneracy of the genetic code, are identical to the DNA sequence of SEQ ID NO: 1 or SEQ ID NO:10, and encode the protein of SEQ ID NO: 2 or SEQ ID NO:11. Further included in the present invention are DNA sequences which hybridize under stringent conditions with the DNA sequence of SEQ ID NO: 1 or SEQ ID NO:10 and encode a protein having BMP-11 activity.

Finally, allelic or other variations of the sequences of SEQ ID NO: 1 or SEQ ID NO: 10, whether such nucleotide changes result in changes in the peptide sequence or not, but where the peptide sequence still has BMP-11 activity, are also included in the present invention.

A further aspect of the invention includes vectors comprising a DNA sequence as described above in operative association with an expression control sequence therefor.

These vectors may be employed in a novel process for producing a BMP-11 protein of the invention in which a cell line transformed with a DNA sequence encoding a BMP-11 protein in operative association with an expression control sequence therefor, is cultured in a suitable culture medium and a BMP-11 protein is recovered and purified therefrom. This process may employ a number of known cells both prokaryotic and eukaryotic as host cells for expression of the polypeptide.

The present invention also includes the use of the DNA sequences and vectors of the invention in gene therapy applications. In such use, the vectors may be transfected into the cells of a patient in vitro, and the cells may be re-introduced into a patient. Alternatively, the vectors may be introduced into a patient in vivo through targeted transfection.

Other aspects and advantages of the present invention will be apparent upon consideration of the following detailed description and preferred embodiments thereof.

Description of the Sequences

SEQ ID NO:1 is a partial nucleotide sequence of the bovine BMP-11 encoding the mature bovine BMP-11 polypeptide.

SEQ ID NO:2 is the amino acid sequence of a partial propeptide and the complete mature bovine BMP-11 polypeptide, encoded by SEQ ID NO:1.

SEQ ID NO:3 is a partial nucleotide sequence of human BMP-11.

SEQ ID NO:4 is a partial amino acid sequence for human BMP-11 polypeptide encoded by SEQ ID NO:3.

SEQ ID NO:5 and 6 are primers to bovine BMP-11 used to isolate the human BMP-11 or other BMP-11 proteins.

SEQ ID NO:7 is a DNA sequence that is inserted into pMT2 CXM to add an XhoI recognition site near the SV40 origin of replication.

SEQ ID NO:8 is a DNA sequence inserted into pMT21 to insert an XhoI recognition site upstream from the DHFR gene.

SEQ ID NO:9 is a DNA sequence comprising a portion of the EMC virus leader sequence.

SEQ ID NO:10 is a DNA sequence encoding a partial propeptide and the complete mature human BMP-11 protein.

SEQ ID NO:11 is the amino acid sequence of a partial propeptide and the complete mature human BMP-11 protein encoded by SEQ ID NO:10.

Detailed Description of the Invention

BMP-11

The bovine BMP-11 nucleotide sequence (SEQ ID NO: 1) and encoded amino acid sequence (SEQ ID NO: 2) and human BMP-11 nucleotide sequence (SEQ ID NO:10) and encoded amino acid sequence (SEQ ID NO:11) are depicted in the Sequence Listings herein. Purified bovine BMP-11 proteins of the present invention are produced by culturing a host cell transformed with a DNA sequence comprising the DNA coding sequence of SEQ ID NO: 1 from nucleotide #375 to #704 or the DNA coding sequence of SEQ ID NO: 10 from nucleotide #760 to #1086 and recovering and purifying from the culture medium a protein which contains the amino acid sequence or a substantially homologous sequence as represented by amino acids # 1 to # 109 of SEQ ID NO: 2 or amino acids 1 to 109 of SEQ ID NO: 11. For production of BMP-11 proteins in mammalian cells, the DNA sequence further comprises a suitable propeptide linked in frame with the above DNA coding sequences for BMP-11. The propeptide may be the native propeptide of BMP-11 or a propeptide from another member of the TGF- β superfamily

The human BMP-11 sequence of the present invention is obtained using the whole or fragments of the bovine BMP-11 DNA sequence, or the partial human BMP-11 sequence of SEQ ID NO: 3

as a probe. Thus, the human BMP-11 DNA sequence comprise the DNA sequence of nucleotides #28 to #185 of SEQ ID NO: 3. The human BMP-11 protein comprise the amino acid sequence of amino acids #1 to #52 of SEQ ID NO: 4.

5 It is expected that BMP-11, as expressed by mammalian cells such as CHO cells, exists as a heterogeneous population of active species of BMP-11 protein with varying N-termini. It is expected that active species will comprise an amino acid sequence beginning at least with the cysteine residue at amino
10 acid #6 of SEQ ID NO:1 or SEQ ID NO:10, or further in the N-terminal direction. Thus, it is expected that DNA sequences encoding active BMP-11 proteins will comprise nucleotides #375 or #390 to 701 of SEQ ID NO:1 or nucleotides #760 or #775 to #1086 of SEQ ID NO:10, and may comprise additional nucleotide
15 sequence in the 5' direction of SEQ ID NO:1 or SEQ ID NO:10.

The N-terminus of human BMP-11 has been experimentally determined by expression in *E. coli* to be as follows: [M]NLGLDXDEHSSE, wherein X designates an amino acid residue with no clear signal, consistent with a location of cysteine at that
20 position. Thus, it is expected that this species of BMP-11 will have an N-terminus at amino acid # 1 of SEQ ID NO:1 or SEQ ID NO: 10, and DNA sequences encoding this species will comprise nucleotides # 375 to # 701 of SEQ ID NO:1 (bovine) or nucleotides # 760 to 1086 of SEQ ID NO:10 (human). The apparent
25 molecular weight of human BMP-11 monomer was determined by SDS-PAGE to be approximately 12 kd. The human BMP-11 protein exists as a clear, colorless solution in 0.1% trifluoroacetic acid.

The BMP-11 proteins recovered from the culture medium are purified by isolating them from other proteinaceous materials
30 from which they are co-produced and from other contaminants present.

BMP-11 proteins may be characterized by the ability to regulate the production of FSH. BMP-11 proteins may further be characterized by the ability to modulate the release of follicle
35 stimulating hormone (FSH) in established *in vitro* bioassays using rat anterior pituitary cells as described [see, e.g., Vale et al, *Endocrinology*, 91:562-572 (1972); Ling et al., *Nature*,

321:779-782 (1986) or Vale et al., *Nature*, 321:776-779 (1986)].

BMP-11 proteins may also be characterized by the ability to induce the formation of bone, cartilage and/or other connective tissue. Such tissue-inducing activity of BMP-11 may further be characterized by the ability to induce the formation of bone, cartilage and/or other connective tissue in the assays described in the examples below.

The BMP-11 proteins provided herein also include factors encoded by the sequences similar to those of SEQ ID NO: 1 or SEQ ID NO:10, but into which modifications are naturally provided (e.g. allelic variations in the nucleotide sequence which may result in amino acid changes in the polypeptide) or deliberately engineered. For example, synthetic polypeptides may wholly or partially duplicate continuous sequences of the amino acid residues of SEQ ID NO:2 or SEQ ID NO:11. These sequences, by virtue of sharing primary, secondary, or tertiary structural and conformational characteristics with inhibin- β polypeptides of SEQ ID NO:2 or SEQ ID NO:11 may possess BMP-11 activity in common therewith. Thus, they may be employed as biologically active substitutes for naturally-occurring BMP-11 polypeptides in therapeutic processes.

Other specific mutations of the sequences of BMP-11 proteins described herein involve modifications of glycosylation sites. These modifications may involve O-linked or N-linked glycosylation sites. For instance, the absence of glycosylation or only partial glycosylation results from amino acid substitution or deletion at asparagine-linked glycosylation recognition sites. The asparagine-linked glycosylation recognition sites comprise tripeptide sequences which are specifically recognized by appropriate cellular glycosylation enzymes. These tripeptide sequences are either asparagine-X-threonine or asparagine-X-serine, where X is usually any amino acid. A variety of amino acid substitutions or deletions at one or both of the first or third amino acid positions of a glycosylation recognition site (and/or amino acid deletion at the second position) results in non-glycosylation at the modified tripeptide sequence. In addition, expression of the

BMP-11 protein in bacterial cells results in non-glycosylated protein, without altering the glycosylation recognition sites.

The present invention also encompasses the novel DNA sequences, free of association with DNA sequences encoding other proteinaceous materials, and coding for expression of BMP-11 proteins. These DNA sequences include those depicted in SEQ ID NO:1 or SEQ ID NO:10 in a 5' to 3' direction and those sequences which hybridize thereto under stringent hybridization conditions, for example 0.1X SSC, 0.1% SDS at 65°C [see, Maniatis et al, Molecular Cloning (A Laboratory Manual), Cold Spring Harbor Laboratory (1982), pages 387 to 389] and encode a protein having BMP-11 activity. These DNA sequences also include those which comprise the DNA sequence of SEQ ID NO:3 and those which hybridize thereto under stringent hybridization conditions and encode a protein having BMP-11 activity.

Similarly, DNA sequences which code for BMP-11 proteins coded for by the sequences of SEQ ID NO:1 or SEQ ID NO:10, but which differ in codon sequence due to the degeneracies of the genetic code or allelic variations (naturally-occurring base changes in the species population which may or may not result in an amino acid change) also encode the novel factors described herein. Variations in the DNA sequences of SEQ ID NO:1 or SEQ ID NO:10 which are caused by point mutations or by induced modifications (including insertion, deletion, and substitution) to enhance the activity, half-life or production of the polypeptides encoded are also encompassed in the invention.

Another aspect of the present invention provides a novel method for producing BMP-11 proteins. The method of the present invention involves culturing a suitable cell line, which has been transformed with a DNA sequence encoding a BMP-11 protein of the invention, under the control of known regulatory sequences. The transformed host cells are cultured and the BMP-11 proteins recovered and purified from the culture medium. The purified proteins are substantially free from other proteins with which they are co-produced as well as from other contaminants.

Suitable cells or cell lines may be mammalian cells, such as Chinese hamster ovary cells (CHO). The selection of suitable

mammalian host cells and methods for transformation, culture, amplification, screening, product production and purification are known in the art. See, e.g., Gething and Sambrook, Nature, 293:620-625 (1981), or alternatively, Kaufman et al, Mol. Cell. Biol., 5(7):1750-1759 (1985) or Howley et al, U.S. Patent 4,419,446. Another suitable mammalian cell line, which is described in the accompanying examples, is the monkey COS-1 cell line. The mammalian cell CV-1 may also be suitable.

Bacterial cells may also be suitable hosts. For example, the various strains of E. coli (e.g., HB101, MC1061) are well-known as host cells in the field of biotechnology. Various strains of B. subtilis, Pseudomonas, other bacilli and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art may also be available as host cells for expression of the polypeptides of the present invention. Additionally, where desired, insect cells may be utilized as host cells in the method of the present invention. See, e.g. Miller et al, Genetic Engineering, 8:277-298 (Plenum Press 1986) and references cited therein.

Another aspect of the present invention provides vectors for use in the method of expression of these novel BMP-11 polypeptides. Preferably the vectors contain the full novel DNA sequences described above which encode the novel factors of the invention. Additionally, the vectors contain appropriate expression control sequences permitting expression of the BMP-11 protein sequences. Alternatively, vectors incorporating modified sequences as described above are also embodiments of the present invention. Additionally, the sequence of SEQ ID NO:1 or SEQ ID NO:10 or other sequences encoding BMP-11 proteins could be manipulated to express a mature BMP-11 by deleting BMP-11 encoding propeptide sequences and replacing them with sequences encoding the complete propeptides of other BMP proteins, activin proteins or other members of the TGF- β superfamily.

The vectors may be employed in the method of transforming cell lines and contain selected regulatory sequences in

operative association with the DNA coding sequences of the invention which are capable of directing the replication and expression thereof in selected host cells. Regulatory sequences for such vectors are known to those skilled in the art and may be selected depending upon the host cells. Such selection is routine and does not form part of the present invention.

For expression in mammalian host cells, the vector may comprise a coding sequence encoding a propeptide suitable for secretion of proteins by the host cell linked in proper reading frame to the coding sequence for mature BMP-11 protein. Suitable propeptide encoding sequences may be obtained from DNA encoding proteins of the TGF- β superfamily of proteins, for example, including BMP-2 through BMP-9. For example, see United States Patent 5,168,150, the disclosure of which is hereby incorporated by reference, in which a DNA encoding a precursor portion of a mammalian protein other than BMP-2 is fused to the DNA encoding a mature BMP-2 protein. Thus, the present invention includes chimeric DNA molecules comprising a DNA sequence encoding a propeptide from a member of the TGF- β superfamily of proteins linked in correct reading frame to a DNA sequence encoding a BMP-11 polypeptide. The term "chimeric" is used to signify that the propeptide originates from a different polypeptide than BMP-11.

A protein of the present invention, which regulates the production of FSH, has possible application in increasing fertility, when expressed in a composition as a homodimer or as a heterodimer with other proteins of the inhibin- β family. The proteins of the present invention may also be useful for contraception, when expressed in a composition as a heterodimer with proteins of the inhibin- α family.

A protein of the present invention, which induces cartilage and/or bone formation in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage defects in humans and other animals. Such a preparation employing a BMP-11 protein may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo

bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery. A BMP-11 protein may be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. BMP-11 polypeptides of the invention may also be useful in the treatment of osteoporosis. A variety of osteogenic, cartilage-inducing and bone inducing factors have been described. See, e.g., European patent applications 148,155 and 169,016 for discussions thereof.

The proteins of the invention may also be used in wound healing and related tissue repair. The types of wounds include, but are not limited to burns, incisions and ulcers. (See, e.g. PCT Publication W084/01106 for discussion of wound healing and related tissue repair).

A further aspect of the invention is a therapeutic method and composition for repairing fractures and other conditions related to cartilage and/or bone defects or periodontal diseases. The invention further comprises therapeutic methods and compositions for wound healing and tissue repair. Such compositions comprise a therapeutically effective amount of at least one of the BMP-11 proteins of the invention in admixture with a pharmaceutically acceptable vehicle, carrier or matrix.

Such a preparation employing a BMP-11 protein may also increase neuronal survival and therefore be useful in transplantation and treatment of conditions exhibiting a decrease in neuronal survival.

It is expected that the BMP-11 proteins of the invention may act in concert with or perhaps synergistically with other related proteins and growth factors. Further therapeutic methods and compositions of the invention therefore comprise a therapeutic amount of at least one BMP-11 protein of the invention with a therapeutic amount of at least one of the BMP proteins or other growth factors disclosed in co-owned patents

and applications described above. Such combinations may comprise separate molecules or heteromolecules comprised of different moieties. For example, a method and composition of the invention may comprise a disulfide linked dimer comprising

5 a BMP-11 protein subunit and a subunit from an inhibin- α protein, an inhibin- β protein or a BMP protein, such as BMP-1 through BMP-10. The agents useful with BMP-11 may include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth

10 factors (TGF- α and TGF- β), and insulin-like growth factor (IGF). Further therapeutic methods and compositions of the invention comprise a therapeutic amount of at least one BMP-11 protein of the invention with a therapeutic amount of at least one of the BMP proteins disclosed in co-owned patents and applications

15 described above. Such combinations may comprise separate molecules of the BMP proteins or heteromolecules comprised of different BMP moieties. For example, a method and composition of the invention may comprise a disulfide linked dimer comprising a BMP-11 protein subunit and a subunit from one of

20 the "BMP" proteins described above. Thus, the present invention includes a purified BMP-11 polypeptide which is a heterodimer wherein one subunit comprises at least the amino acid sequence from amino acid #1 to amino acid #109 of SEQ ID NO:2 or SEQ ID NO:11, and one subunit comprises an amino acid sequence for a

25 bone morphogenetic protein selected from the group consisting of BMP-1, BMP-2, BMP-3, BMP-4, BMP-5, BMP-6, BMP-7, BMP-8 and BMP-9. A further embodiment may comprise a heterodimer of BMP-11 moieties. Further, BMP-11 proteins may be combined with other agents beneficial to the treatment of the bone and/or cartilage

30 defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth factor (EGF), fibroblast growth factor (FGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- α and TGF- β), and k-fibroblast growth factor (kFGF), parathyroid hormone (PTH),

35 leukemia inhibitory factor (LIF/HILDA/DIA), insulin-like growth factors (IGF-I and IGF-II). Portions of these agents may also be used in compositions of the present invention.

The BMP-11 proteins of the present invention may also be used in compositions combined with bone morphogenetic proteins. See for example, Ogawa et al., WO 92/14481 (1992); Ogawa et al.,
5 *J. Biol. Chem.*, 267:14233-14237 (1992). The bone morphogenetic proteins useful in such compositions include BMP-1, BMP-2, BMP-3, BMP-4, BMP-5, BMP-6 and BMP-7, disclosed for instance in United States Patents 5,108,922; 5,013,649; 5,116,738; 5,106,748; 5,187,076; and 5,141,905; BMP-8, disclosed in PCT
10 publication WO91/18098; and BMP-9, disclosed in PCT publication WO93/00432; and BMP-10 disclosed in co-pending patent application serial number 08/061,695, filed on May 12, 1993.

The preparation and formulation of such physiologically acceptable protein compositions, having due regard to pH,
15 isotonicity, stability and the like, is within the skill of the art. The therapeutic compositions are also presently valuable for veterinary applications due to the lack of species specificity in BMP and TGF proteins. Particularly domestic animals and thoroughbred horses in addition to humans are
20 desired patients for such treatment with BMP-11 of the present invention.

The therapeutic method includes administering the composition topically, systemically, or locally as an implant or device. When administered, the therapeutic composition for use
25 in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and
30 tissue repair. Therapeutically useful agents other than the BMP-11 proteins which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the BMP-11 composition in the methods of the invention.

35 Preferably for bone, cartilage or other connective tissue formation, the composition includes a matrix capable of delivering BMP-11 or other BMP proteins to the site of tissue

damage in need of repair, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. The matrix may provide slow release of BMP-11 and/or other bone inductive protein, as well as proper presentation and appropriate environment for cellular infiltration. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the BMP-11 compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid and polyanhydrides. Other potential materials are biodegradable and biologically well defined, such as bone, tendon or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

Progress can be monitored by periodic assessment of bone growth and/or repair. The progress can be monitored, for example, x-rays, histomorphometric determinations and tetracycline labeling.

The dosage regimen will be determined by the attending physician considering various factors which modify the action of the BMP-11 protein, e.g. the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of BMP protein or growth factor present in the composition. The dosage may also vary with the type of matrix used.

The following examples illustrate practice of the present invention in recovering and characterizing bovine BMP-11 protein and employing it to recover the human and other BMP-11 proteins, obtaining the human proteins and expressing the proteins via recombinant techniques.

EXAMPLE 1

Bovine BMP-11

800,000 recombinants of a bovine genomic library constructed in the vector λ EMBL3 are plated at a density of 8000 recombinant bacteriophage plaques per plate on 100 plates. Duplicate nitrocellulose replicas of the recombinant bacteriophage plaques are made from these plates and amplified. A fragment of human BMP-7 DNA corresponding to nucleotides #1081 to #1403 (Figure 4, United States Patent 5,141,905) is ^{32}P -labelled by the random priming procedure of Feinberg et al. [Anal. Biochem. 132: 6-13 (1983)] and hybridized to one set of filters in standard hybridization buffer (SHB) (5x SSC, 0.170 SDS, 5x Denhardt's, 100 $\mu\text{g}/\text{ml}$ Salmon sperm DNA) at 60°C for 2 to 3 days. The filters are washed under reduced stringency conditions (4X SSC, 0.1% SDS at 60°C). Multiple positively hybridizing recombinants are noted. 52 positively hybridizing recombinant bacteriophage plaques are selected and replated for secondaries. Duplicate nitrocellulose replicas of the recombinant plaques are made from these 52 secondary plates and amplified. One set of nitrocellulose filters is hybridized to the human BMP-7 DNA probe as described above and washed under the same reduced stringency conditions. The other set of filters is hybridized to a mixed BMP-5, BMP-6, and BMP-7 probe in SHB at 65°C overnight and washed with a 0.1X SSC, 0.1% SDS at 65°C (stringent hybridization and wash conditions). The mixed probe consists of relatively equal amounts of ^{32}P -labelled DNA fragments comprising nucleotides #1452 to #2060 (Figure 4, United States Patent 5,106,748) of the human BMP-5 sequence, nucleotides #1395 to #1698 (Figure 4, United States Patent 5,187,076) of the human BMP-6 sequence, and nucleotides #1081 to #1403 (Figure 4, United States Patent 5,141,905) of the human BMP-7 sequence. The BMP-5, BMP-6 and BMP-7 DNA fragments are

³²P-labelled by the random priming procedure and equal numbers of counts per minute (cpms) of each probe are combined and added to the SHB containing the other set of nitrocellulose filter replicas of the 52 secondary plates. 14 recombinants, which hybridized positively to the human BMP-7 probe under the reduced stringency conditions and exhibited weak or no hybridization to the mixed BMP-5/6/7 probe under high stringency conditions, are selected for further analysis. All 14 recombinants which exhibit these hybridization characteristics are plaque purified and bacteriophage DNA is prepared from each. The positively hybridizing region of one of the 14 recombinants exhibiting the hybridization characteristics described above, designated λ 7r-30, is localized to a 0.5 kb SacI restriction fragment. This fragment is subcloned into a plasmid vector (pGEM-3) and DNA sequence analysis is performed. The partial DNA sequence (SEQUENCE ID NO. 1) and derived amino acid sequence (SEQUENCE ID NO. 2) of clone λ 7r-30 are shown in the Sequence Listings.

The bacteriophage λ 7r-30 has been deposited with the ATCC on April 7, 1993, and accorded the accession number ATCC 75439. This deposit meets the requirements of the Budapest Treaty of the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedures and regulations thereunder.

This λ 7r-30 clone encodes at least a portion of the bovine BMP-11 protein of the present invention. The nucleotide sequence of clone λ 7r-30 contains an open reading frame of 456 nucleotides #246-701 of SEQ ID NO:1. The nucleotide sequence of #324 to #701 of SEQ ID NO:1 defines an open reading frame of 378 nucleotides, encoding at least 126 amino acids of the C-terminal portion of a bovine BMP-11 protein, as determined by alignment to other BMP proteins and other proteins within the TGF- β family. The nucleotide sequence #246 to #323 defines an open reading frame contiguous with the sequence encoding the predicted 126 amino acid BMP-11 peptide, however a reduced degree of amino acid identity of the peptide deduced from this region of DNA sequence (#246 to #323) to other BMP proteins and other proteins of the TGF- β family and the presence of multiple potential splice acceptor consensus sequences make it difficult

to define the 5' limit of this exon of the bovine BMP-11 gene. The presence of an in-frame stop codon at nucleotide positions #243 to #245 indicates that nucleotide sequence of clone λ 7r-30 contains at least one exon/intron boundary of the bovine BMP-11 gene.

Based upon the knowledge of other proteins within the TGF- β family, it is predicted that the BMP-11 precursor polypeptide would be cleaved at the multibasic sequence ARG-SER-ARG-ARG in agreement with a proposed consensus proteolytic processing sequence of ARG-X-X-ARG. Cleavage of the BMP-11 precursor polypeptide is expected to generate a 109 amino acid mature peptide beginning with the amino acid ASN at position #1. The processing of BMP-11 into the mature form is expected to involve dimerization and removal of the N-terminal region in a manner analogous to the processing of the related protein TGF- β [Gentry et al., Molec. & Cell. Biol., 8:4162 (1988); Derynck et al., Nature, 316:701(1985)].

It is contemplated therefore that the mature active species of BMP-11 comprises a homodimer of two polypeptide subunits, each subunit comprising amino acids # 1 to # 109 with a predicted molecular weight of approximately 12,000 daltons. Further active species are contemplated comprising amino acids #6 to #109, thereby including the first conserved cysteine residue. As with other members of the TGF- β family of proteins, the carboxy-terminal region of the BMP-11 protein exhibits greater sequence conservation than the more amino-terminal portion. The percent amino acid identity of the BMP-11 protein in the cysteine-rich C-terminal domain (amino acids #6 to #109) to the corresponding region of other proteins within the TGF- β family is as follows: BMP-2, 39%; BMP-3, 37%; BMP-4, 37%; BMP-5, 42%; BMP-6, 45%; BMP-7, 42%; BMP-8, 39%; BMP-9, 40%; Vg1, 39%; GDF-1, 34%; TGF- β 1, 36%; TGF- β 2, 38%; TGF- β 3, 38%; inhibin β (B), 41%; inhibin β (A), 39%.

EXAMPLE 2

Human BMP-11

Bovine and human BMP-11 genes are presumed to be

significantly homologous, therefore the bovine coding sequence or a portion thereof is used as a probe to screen a human genomic library or as a probe to identify a human cell line or tissue which synthesizes the analogous human protein. A human
5 genomic library, such as Stratagene catalog #944201, may be screened with such a probe, and presumptive positives isolated and DNA sequence obtained. Evidence that this recombinant encodes a portion of the human BMP-11 relies on the bovine/human protein and gene structure homologies.

10 Once a recombinant bacteriophage containing DNA encoding a portion of the human BMP-11 molecule is obtained, the human coding sequence can be used as a probe to identify a human cell line or tissue which synthesizes BMP-11 mRNA. Alternatively, the bovine BMP-11 coding sequence can be used as a probe to
15 identify such human cell line or tissue. Briefly described, RNA is extracted from a selected cell or tissue source and either electrophoresed on a formaldehyde agarose gel and transferred to nitrocellulose, or reacted with formaldehyde and spotted on nitrocellulose directly. The nitrocellulose is then hybridized
20 to a probe derived from a coding sequence of the bovine or human BMP-11. Alternatively, the bovine BMP-11 coding sequence is used to design oligonucleotide primers which will specifically amplify a portion of the BMP-11 encoding sequence located in the region located between the primers utilized to perform the
25 specific amplification reaction. It is contemplated that bovine and human BMP-11 sequences would allow one to specifically amplify corresponding human BMP-11 encoding sequences from mRNA, cDNA or genomic DNA templates. Once a positive source has been identified by one of the above described methods, mRNA is
30 selected by oligo (dT) cellulose chromatography and cDNA is synthesized and cloned in λ gt10 or other λ bacteriophage vectors known to those skilled in the art. (i.e. λ ZAP) by established techniques (Toole et al., supra). It is also possible to perform the oligonucleotide primer directed amplification
35 reaction, described above, directly on a pre-established human cDNA or genomic library which has been cloned into a λ bacteriophage vector. In such cases, a library which yields a

specifically amplified DNA product encoding a portion of human BMP-11 protein could be screened directly, utilizing the fragment of amplified BMP-11 encoding DNA as a probe.

Oligonucleotide primers designed on the basis of the DNA sequence of the bovine BMP-11 genomic clone λ 7r-30 are predicted to allow the specific amplification of human BMP-11 encoding sequences. The following oligonucleotide primer is designed on the basis of nucleotides #501 to #521 of the DNA sequence set forth in SEQ ID NO. 1 and synthesized on an automated DNA synthesizer.

Primer C: TAGTCTAGATGCTCCGGCCAGTGCGAGTAC

The first nine nucleotides of primer C (underlined) comprise the recognition sequence for the restriction endonuclease XbaI which can be utilized to facilitate the manipulation of a specifically amplified DNA sequence encoding the BMP-11 protein of the invention and are thus not derived from the DNA sequence presented in SEQ ID NO: 1.

The following oligonucleotide primer is designed on the basis of nucleotides # 701 to # 678 of the DNA sequence set forth in SEQ ID NO. 1 and synthesized on an automated DNA synthesizer:

Primer D: TGCGGATCCGGAGCAGCCACAGCGATCCAC

The first nine nucleotides of primer D (underlined) comprise the recognition sequence for the restriction endonuclease BamHI which can be utilized to facilitate the manipulation of a specifically amplified DNA sequence encoding the BMP-11 protein of the invention and are thus not derived from the DNA sequence present in SEQ ID NO:1.

The standard nucleotide symbols in the above identified primers are as follows: A, adenosine; C, cytosine, G, guanine; and T, thymine.

Primers C and D identified above are utilized as primers to allow the amplification of a specific nucleotide from human genomic DNA. The amplification reaction is performed as follows:

Human genomic DNA (source: peripheral blood lymphocytes) is denatured at 100°C for five minutes and then chilled on ice

prior to addition to a reaction mixture containing 200 μ M each deoxynucleotide triphosphates (dATP, dGTP, dCTP and dTTP) 10 mM Tris-HCl pH8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.001% gelatin, 1.25 units Taq DNA polymerase, 100 pM oligonucleotide primer C and 100 pM oligonucleotide primer D. This reaction mixture is then subjected to thermal cycling in the following manner: 3 minutes at 94°C, 1 minute at 50°C, 1 minute at 72°C for one cycle, then 1 minute at 94°C, 1 minute at 50°C, 1 minute at 72°C for thirty-nine cycles.

The DNA which is specifically amplified by this reaction is separated from the excess oligonucleotide primers C and D utilized to initiate the amplification by the use of a DNA purification resin based protocol under the conditions suggested by the manufacturer. The resulting DNA product is digested with the restriction endonucleases XbaI and BamHI, phenol extracted, chloroform extracted. Buffer exchange and removal of small fragments of DNA resulting from the XbaI/BamHI restriction digest is accomplished by dilution of the digested DNA product in 10 mM Tris-HCl pH8.0, 1 mM EDTA followed by centrifugation through a centriconTM 30 microconcentrator (W.R. Grace & Co., Beverly, Ma.; Product #4209). The resulting XbaI/BamHI digested amplified DNA product is subcloned into a plasmid vector (pBluescript) between the XbaI and BamHI restriction sites of the polylinker region. DNA sequence analysis of the resulting subclones indicates that the specifically amplified DNA sequence product encodes a portion of the human BMP-11 protein of this invention. The DNA sequence (SEQ ID NO:3) and derived amino acid sequence (SEQ ID NO:4) of this specifically amplified DNA fragment are set forth in the Sequence Listings.

Nucleotides #1 to #27 of this sequence comprise a portion of oligonucleotide primer C and nucleotides #186 to #213 comprise a portion of oligonucleotide primer D utilized to perform the specific amplification reaction. Due to the function of oligonucleotide primers C and D (designed on the basis of bovine BMP-11 DNA sequence) in initiating the amplification reaction, they may not correspond exactly to the actual sequence encoding a human BMP-11 and are therefore not

translated in the above amino acid sequence derivation. The DNA sequence, from nucleotide #28 to #185 of SEQ ID NO: 3, or portions thereof, specifically amplified from the human genomic DNA template can be utilized as a probe to identify additional human BMP-11 encoding sequences from human genomic or human cDNA libraries by standard hybridization/ screening techniques known to those skilled in the art.

One million, two hundred thousand recombinants of a human fetal brain cDNA library (Stratagene catalog # 936206) constructed in the vector λ ZAPII are plated at a density of 24,000 recombinant bacteriophage plaques per plate on 50 plates. Duplicate nitrocellulose replicas of the recombinant bacteriophage plaques are made from these plates. An oligonucleotide probe designed on the basis of nucleotides #53-#82 of SEQ ID NO:3 is synthesized on an automated DNA synthesizer. This oligonucleotide probe is radioactively labelled with $\gamma^{32}\text{P}$ -ATP and is hybridized to both sets of the duplicate nitrocellulose replicas in SHB at 65°C. Nine positively hybridizing recombinants are noted. One of the positively hybridizing recombinants, named λ FB30.5, is plaque purified. Bacteriophage plate stocks of the purified λ FB30.5 cDNA clone are prepared and bacteriophage DNA is isolated. A bacterial plasmid named FB30.5, generated by the in vivo excision protocol described by the supplier (Stratagene) and containing the entire insert of the λ FB30.5 bacteriophage cDNA clone, has been deposited with the ATCC, 12301 Parklawn Drive, Rockville, Maryland USA under the requirements of the Budapest Treaty and designated as ATCC # _____. A portion of the DNA sequence of clone FB30.5 is set forth in SEQ ID NO: 10.

One million recombinants of a human genomic library (Stratagene Catalog # 944201) constructed in the vector λ FIX are plated at a density of 20,000 recombinant bacteriophage plaques per plate on 50 plates. Duplicate nitrocellulose replicas of the recombinant bacteriophage plaques are made from these plates. An oligonucleotide probe designed on the basis of nucleotides #57-#86 of SEQ ID NO:10, with the exception of an inadvertent substitution of CAC for GCG at nucleotides #59-#61

of SEQ ID NO:10, is synthesized on an automated DNA synthesizer. This oligonucleotide probe is radioactively labelled with $\gamma^{32}\text{P}$ -ATP and is hybridized to both sets of the duplicate nitrocellulose replicas in SHB at 65°C. Five positively hybridizing recombinants are noted. One of the positively hybridizing recombinants, named 30GEN.4, is plaque purified. Bacteriophage plate stocks of the purified 30GEN.4 genomic clone are prepared and bacteriophage DNA is isolated. A bacteriophage stock of this genomic clone has been deposited with the ATCC, 12301 Parklawn Drive, Rockville, Maryland USA under the requirements of the Budapest Treaty and designated as ATCC # _____. A portion of the DNA sequence of clone 30GEN.4 is set forth in SEQ ID NO: 10. A portion of the DNA sequence of the genomic clone 30GEN.4 was determined to be identical to a portion of the DNA sequence of the cDNA clone FB30.5. The extent of this overlap (nucleotides #1-#198) of SEQ ID NO:10 were used as a basis to compile the partial coding sequence of the BMP-10 protein. The genomic clone 30GEN.4 is expected to contain additional 5' coding sequences of the human BMP-11 protein which are expected to encode the remainder of the BMP-11 precursor polypeptide, including the initiator methionine. The partial sequence of human BMP-11 is presented in SEQ ID NO:10 and it should be noted that nucleotides #1-198 have been determined to be present in both the 30GEN.4 genomic clone and the FB30.5 cDNA clone while nucleotides #199-#1270 are derived entirely from the cDNA clone FB30.5. SEQ ID NO:10 predicts a human BMP11 precursor protein of at least 362 amino acids. Based on the knowledge of other BMPs and other proteins within the TGF- β family, it is predicted that the precursor polypeptide would be cleaved at the multibasic sequence ARG-SER-ARG-ARG (amino acids #-4 through #-1 of SEQ ID NO:11) in agreement with the proposed consensus proteolytic processing sequence ARG-X-X-ARG. Cleavage of the human BMP-11 precursor polypeptide at this location would generate a 109 amino acid mature peptide beginning with the amino acid ASN at position #1 of SEQ ID NO:11. The processing of human BMP-11 into the mature form is expected to involve dimerization and removal of the N-terminal

region in a manner analogous to the processing of the related protein TGF- β [L.E. Gentry, et al. Molec. & Cell. Biol. 8:4162 (1988); R.Derynck, et al., Nature 316:701 (1985)]. It is contemplated that the mature active species of human BMP-11 comprises a homodimer of two polypeptide subunits, each subunit comprising amino acids #1-#108 of SEQ ID NO:11, with a predicted molecular weight of 12,000 daltons. Further active species are contemplated comprising amino acids #7-#108 of SEQ ID NO:11, thereby including the first conserved cysteine residue. Heterodimeric molecules comprising one subunit of BMP-11 and another subunit of another member of the BMP/TGF- β superfamily are also contemplated.

EXAMPLE 3

Expression of BMP-11

In order to produce bovine, human or other mammalian BMP-11 proteins, the DNA encoding it is transferred into an appropriate expression vector and introduced into mammalian cells or other preferred eukaryotic or prokaryotic hosts by conventional genetic engineering techniques. The preferred expression system for biologically active recombinant human BMP-11 is contemplated to be stably transformed mammalian cells.

One skilled in the art can construct mammalian expression vectors by employing the sequence of SEQ ID NO:1 or SEQ ID NO:10, or other DNA sequences encoding BMP-11 proteins or other modified sequences and known vectors, such as pCD [Okayama et al., Mol. Cell Biol., 2:161-170 (1982)], pJL3, pJL4 [Gough et al., EMBO J., 4:645-653 (1985)] and pMT2 CXM.

The mammalian expression vector pMT2 CXM is a derivative of p91023(b) (Wong et al., Science 228:810-815, 1985) differing from the latter in that it contains the ampicillin resistance gene in place of the tetracycline resistance gene and further contains a XhoI site for insertion of cDNA clones. The functional elements of pMT2 CXM have been described (Kaufman, R.J., 1985, Proc. Natl. Acad. Sci. USA 82:689-693) and include the adenovirus VA genes, the SV40 origin of replication including the 72 bp enhancer, the adenovirus major late promoter including a 5' splice site and the majority of the adenovirus

tripartite leader sequence present on adenovirus late mRNAs, a 3' splice acceptor site, a DHFR insert, the SV40 early polyadenylation site (SV40), and pBR322 sequences needed for propagation in E. coli.

5 Plasmid pMT2 CXM is obtained by EcoRI digestion of pMT2-VWF, which has been deposited with the American Type Culture Collection (ATCC), Rockville, MD (USA) under accession number ATCC 67122. EcoRI digestion excises the cDNA insert present in pMT2-VWF, yielding pMT2 in linear form which can be ligated and
10 used to transform E. coli HB 101 or DH-5 to ampicillin resistance. Plasmid pMT2 DNA can be prepared by conventional methods. pMT2 CXM is then constructed using loopout/in mutagenesis [Morinaga, et al., Biotechnology 84: 636 (1984). This removes bases 1075 to 1145 relative to the Hind III site
15 near the SV40 origin of replication and enhancer sequences of pMT2. In addition it inserts the following sequence:

5' PO-CATGGGCAGCTCGAG-3'

at nucleotide 1145. This sequence contains the recognition site for the restriction endonuclease Xho I. A derivative of
20 pMT2CXM, termed pMT23, contains recognition sites for the restriction endonucleases PstI, Eco RI, SalI and XhoI. Plasmid pMT2 CXM and pMT23 DNA may be prepared by conventional methods.

pEMC281 derived from pMT21 may also be suitable in practice of the invention. pMT21 is derived from pMT2 which is derived
25 from pMT2-VWF. As described above EcoRI digestion excises the cDNA insert present in pMT-VWF, yielding pMT2 in linear form which can be ligated and used to transform E. Coli HB 101 or DH-5 to ampicillin resistance. Plasmid pMT2 DNA can be prepared by conventional methods.

30 pMT21 is derived from pMT2 through the following two modifications. First, 76 bp of the 5' untranslated region of the DHFR cDNA including a stretch of 19 G residues from G/C tailing for cDNA cloning is deleted. In this process, a XhoI site is inserted to obtain the following sequence immediately
35 upstream from DHFR: 5' -CTGCAGGCGAGCCTGAATTCCTCGAGCCATCATG-3'
PstI Eco RI XhoI

Second, a unique ClaI site is introduced by digestion with

EcoRV and XbaI, treatment with Klenow fragment of DNA polymerase I, and ligation to a ClaI linker (CATCGATG). This deletes a 250 bp segment from the adenovirus associated RNA (VAI) region but does not interfere with VAI RNA gene expression or function. pMT21 is digested with EcoRI and XhoI, and used to derive the vector pEMC2B1.

A portion of the EMCV leader is obtained from pMT2-ECAT1 [S.K. Jung, et al, *J. Virol* 63:1651-1660 (1989)] by digestion with Eco RI and PstI, resulting in a 2752 bp fragment. This fragment is digested with TaqI yielding an Eco RI-TaqI fragment of 508 bp which is purified by electrophoresis on low melting agarose gel. A 68 bp adapter and its complementary strand are synthesized with a 5' TaqI protruding end and a 3' XhoI protruding end which has the following sequence:

5'-CGAGGTTAAAAACGTCTAGGCCCCCCGAACCACGGGGACGTGGTTCCTTT
TaqI

GAAAAACACGATTGC-3'
XhoI

This sequence matches the EMC virus leader sequence from nucleotide 763 to 827. It also changes the ATG at position 10 within the EMC virus leader to an ATT and is followed by a XhoI site. A three way ligation of the pMT21 Eco RI-XhoI fragment, the EMC virus EcoRI-TaqI fragment, and the 68 bp oligonucleotide adapter TaqI-XhoI adapter resulting in the vector pEMC2β1.

This vector contains the SV40 origin of replication and enhancer, the adenovirus major late promoter, a cDNA copy of the majority of the adenovirus tripartite leader sequence, a small hybrid intervening sequence, an SV40 polyadenylation signal and the adenovirus VA I gene, DHFR and β-lactamase markers and an EMC sequence, in appropriate relationships to direct the high level expression of the desired cDNA in mammalian cells.

The construction of vectors may involve modification of the BMP-11 DNA sequences. For instance, BMP-11 cDNA can be modified by removing the non-coding nucleotides on the 5' and 3' ends of the coding region. The deleted non-coding nucleotides may or

may not be replaced by other sequences known to be beneficial for expression. These vectors are transformed into appropriate host cells for expression of BMP-11 proteins. Additionally, the sequence of SEQ ID NO:1 or SEQ ID NO:10 or other sequences encoding BMP-11 proteins could be manipulated to express a mature BMP-11 by deleting BMP-11 encoding propeptide sequences and replacing them with sequences encoding the complete propeptides of other BMP proteins, activin proteins or other members of the TGF- β superfamily.

One skilled in the art can manipulate the sequences of SEQ ID NO:1 or SEQ ID NO:10 by eliminating or replacing the mammalian regulatory sequences flanking the coding sequence with bacterial sequences to create bacterial vectors for intracellular or extracellular expression by bacterial cells. For example, the coding sequences could be further manipulated (e.g. ligated to other known linkers or modified by deleting non-coding sequences therefrom or altering nucleotides therein by other known techniques). The modified BMP-11 coding sequence could then be inserted into a known bacterial vector using procedures such as described in T. Taniguchi et al., Proc. Natl Acad. Sci. USA, 77:5230-5233 (1980). This exemplary bacterial vector could then be transformed into bacterial host cells and a BMP-11 protein expressed thereby. For a strategy for producing extracellular expression of BMP-11 proteins in bacterial cells, see, e.g. European patent application EPA 177,343.

Similar manipulations can be performed for the construction of an insect vector [See, e.g. procedures described in published European patent application 155,476] for expression in insect cells. A yeast vector could also be constructed employing yeast regulatory sequences for intracellular or extracellular expression of the factors of the present invention by yeast cells. [See, e.g., procedures described in published PCT application WO86/00639 and European patent application EPA 123,289].

A method for producing high levels of a BMP-11 protein of the invention in mammalian cells may involve the construction of

cells containing multiple copies of the heterologous BMP-11 gene. The heterologous gene is linked to an amplifiable marker, e.g. the dihydrofolate reductase (DHFR) gene for which cells containing increased gene copies can be selected for propagation in increasing concentrations of methotrexate (MTX) according to the procedures of Kaufman and Sharp, J. Mol. Biol., 159:601-629 (1982). This approach can be employed with a number of different cell types.

For example, a plasmid containing a DNA sequence for a BMP-11 of the invention in operative association with other plasmid sequences enabling expression thereof and the DHFR expression plasmid pAdA26SV(A)3 [Kaufman and Sharp, Mol. Cell. Biol., 2:1304 (1982)] can be co-introduced into DHFR-deficient CHO cells, DUKX-BII, by various methods including calcium phosphate coprecipitation and transfection, electroporation or protoplast fusion. DHFR expressing transformants are selected for growth in alpha media with dialyzed fetal calf serum, and subsequently selected for amplification by growth in increasing concentrations of MTX (e.g. sequential steps in 0.02, 0.2, 1.0 and 5uM MTX) as described in Kaufman et al., Mol Cell Biol., 5:1750 (1983). Transformants are cloned, and biologically active BMP-11 expression is monitored by one or more of the BMP-11 activity assays described in Examples 5 to 8 below. BMP-11 expression should increase with increasing levels of MTX resistance. BMP-11 polypeptides are characterized using standard techniques known in the art such as pulse labeling with [35S] methionine or cysteine and polyacrylamide gel electrophoresis. Similar procedures can be followed to produce other related BMP-11 proteins.

EXAMPLE 4

Biological Activity of Expressed BMP-11

To measure the biological activity of the expressed BMP-11 proteins obtained in Example 3 above, the proteins are recovered from the cell culture and purified by isolating the BMP-11 proteins from other proteinaceous materials with which they are co-produced as well as from other contaminants. The purified protein may be assayed in accordance with the assays for BMP-11

activity described in Examples 5 to 8 below.

EXAMPLE 5

W-20 BIOASSAYS

A. Description of W-20 cells

5 Use of the W-20 bone marrow stromal cells as an indicator cell line is based upon the conversion of these cells to osteoblast-like cells after treatment with a BMP protein [Thies et al, Journal of Bone and Mineral Research, 5:305 (1990); and Thies et al, Endocrinology, 130:1318 (1992)]. Specifically, W-
10 20 cells are a clonal bone marrow stromal cell line derived from adult mice by researchers in the laboratory of Dr. D. Nathan, Children's Hospital, Boston, MA. Treatment of W-20 cells with certain BMP proteins results in (1) increased alkaline phosphatase production, (2) induction of PTH stimulated cAMP,
15 and (3) induction of osteocalcin synthesis by the cells. While (1) and (2) represent characteristics associated with the osteoblast phenotype, the ability to synthesize osteocalcin is a phenotypic property only displayed by mature osteoblasts. Furthermore, to date we have observed conversion of W-20 stromal
20 cells to osteoblast-like cells only upon treatment with BMPs. In this manner, the in vitro activities displayed by BMP treated W-20 cells correlate with the in vivo bone forming activity known for BMPs.

25 Below two in vitro assays useful in comparison of BMP activities of novel osteoinductive molecules are described.

B. W-20 Alkaline Phosphatase Assay Protocol

30 W-20 cells are plated into 96 well tissue culture plates at a density of 10,000 cells per well in 200 μ l of media (DME with 10% heat inactivated fetal calf serum, 2 mM glutamine and 100 Units/ml penicillin + 100 μ g/ml streptomycin. The cells are allowed to attach overnight in a 95% air, 5% CO₂ incubator at 37°C.

35 The 200 μ l of media is removed from each well with a multichannel pipettor and replaced with an equal volume of test sample delivered in DME with 10% heat inactivated fetal calf serum, 2 mM glutamine and 1% penicillin-streptomycin. Test substances are assayed in triplicate.

The test samples and standards are allowed a 24 hour incubation period with the W-20 indicator cells. After the 24 hours, plates are removed from the 37°C incubator and the test media are removed from the cells.

5 The W-20 cell layers are washed 3 times with 200 µl per well of calcium/magnesium free phosphate buffered saline and these washes are discarded.

10 50 µl of glass distilled water is added to each well and the assay plates are then placed on a dry ice/ethanol bath for quick freezing. Once frozen, the assay plates are removed from the dry ice/ethanol bath and thawed at 37°C. This step is repeated 2 more times for a total of 3 freeze-thaw procedures. Once complete, the membrane bound alkaline phosphatase is available for measurement.

15 50 µl of assay mix (50 mM glycine, 0.05% Triton X-100, 4 mM MgCl₂, 5 mM p-nitrophenol phosphate, pH = 10.3) is added to each assay well and the assay plates are then incubated for 30 minutes at 37°C in a shaking waterbath at 60 oscillations per minute.

20 At the end of the 30 minute incubation, the reaction is stopped by adding 100 µl of 0.2 N NaOH to each well and placing the assay plates on ice.

25 The spectrophotometric absorbance for each well is read at a wavelength of 405 nanometers. These values are then compared to known standards to give an estimate of the alkaline phosphatase activity in each sample. For example, using known amounts of p-nitrophenol phosphate, absorbance values are generated. This is shown in Table I.

Table I

Absorbance Values for Known Standards of P-Nitrophenol Phosphate	
<u>P-nitrophenol phosphate umoles</u>	<u>Mean absorbance (405 nm)</u>
0.000	0
0.006	0.261 +/- .024
0.012	0.521 +/- .031
0.018	0.797 +/- .063
0.024	1.074 +/- .061
0.030	1.305 +/- .083

5

Absorbance values for known amounts of BMPs can be determined and converted to μ moles of p-nitrophenol phosphate cleaved per unit time as shown in Table II.

Table II

Alkaline Phosphatase Values for W-20 Cells Treating with BMP-2		
<u>BMP-2 concentration ng/ml</u>	<u>Absorbance Reading 405 nmeters</u>	<u>umoles substrate per hour</u>
0	0.645	0.024
1.56	0.696	0.026
3.12	0.765	0.029
6.25	0.923	0.036
12.50	1.121	0.044
25.0	1.457	0.058
50.0	1.662	0.067
100.0	1.977	0.080

10

These values are then used to compare the activities of known amounts of BMP-11 to BMP-2.

C. Osteocalcin RIA Protocol

W-20 cells are plated at 10^6 cells per well in 24 well

multiwell tissue culture dishes in 2 mls of DME containing 10% heat inactivated fetal calf serum, 2 mM glutamine. The cells are allowed to attach overnight in an atmosphere of 95% air 5% CO₂ at 37°C.

5 The next day the medium is changed to DME containing 10% fetal calf serum, 2 mM glutamine and the test substance in a total volume of 2 ml. Each test substance is administered to triplicate wells. The test substances are incubated with the W-20 cells for a total of 96 hours with replacement at 48 hours by
10 the same test medias.

At the end of 96 hours, 50 µl of the test media is removed from each well and assayed for osteocalcin production using a radioimmunoassay for mouse osteocalcin. The details of the assay are described in the kit manufactured by Biomedical
15 Technologies Inc., 378 Page Street, Stoughton, MA 02072. Reagents for the assay are found as product numbers BT-431 (mouse osteocalcin standard), BT-432 (Goat anti-mouse Osteocalcin), BT-431R (iodinated mouse osteocalcin), BT-415 (normal goat serum) and BT-414 (donkey anti goat IgG). The RIA
20 for osteocalcin synthesized by W-20 cells in response to BMP treatment is carried out as described in the protocol provided by the manufacturer.

The values obtained for the test samples are compared to values for known standards of mouse osteocalcin and to the
25 amount of osteocalcin produced by W-20 cells in response to challenge with known amounts of BMP-2. The values for BMP-2 induced osteocalcin synthesis by W-20 cells is shown in Table III.

Table III

Osteocalcin Synthesis by W-20 Cells	
<u>BMP-2 Concentration ng/ml</u>	<u>Osteocalcin Synthesis ng/well</u>
0	0.8
2	0.9
4	0.8
8	2.2
16	2.7
31	3.2
62	5.1
125	6.5
250	8.2
500	9.4
1000	10.0

EXAMPLE 6**Rosen-Modified Sampath-Reddi Assay.**

5 A modified version of the rat bone formation assay described in Sampath and Reddi, Proc. Natl. Acad. Sci. USA,
80:6591-6595 (1983) is used to evaluate bone, cartilage and/or
other connective tissue inductive activity of BMP-11 proteins. This modified assay is herein called the Rosen-modified Sampath-
Reddi assay. The ethanol precipitation step of the Sampath-
10 Reddi procedure is replaced by dialyzing (if the composition is a solution) or diafiltering (if the composition is a suspension)
the fraction to be assayed against water. The solution or
suspension is then equilibrated to 0.1% TFA. The resulting
solution is added to 20 mg of rat matrix. A mock rat matrix
sample not treated with the protein serves as a control. This
15 material is frozen and lyophilized and the resulting powder
enclosed in #5 gelatin capsules. The capsules are implanted
subcutaneously in the abdominal thoracic area of 21-49 day old
male Long Evans rats. The implants are removed after 7-14 days.
Half of each implant is used for alkaline phosphatase analysis
20 [see, Reddi et al, Proc. Natl. Acad. Sci., 69:1601 (1972)].

The other half of each implant is fixed and processed for
histological analysis. 1 μ m glycolmethacrylate sections are
stained with Von Kossa and acid fuchsin to score the amount of
induced bone and cartilage formation present in each implant.
25 The terms +1 through +5 represent the area of each histological
section of an implant occupied by new bone and/or cartilage
cells and matrix. A score of +5 indicates that greater than 50%
of the implant is new bone and/or cartilage produced as a direct
result of protein in the implant. A score of +4, +3, +2, and +1
30 would indicate that greater than 40%, 30%, 20% and 10%
respectively of the implant contains new cartilage and/or bone.

The BMP-11 proteins of this invention may be assessed for
activity on this assay.

EXAMPLE 7**Biological Activity of Expressed BMP-11**

To measure the biological activity of the expressed BMP-11 proteins obtained in Example 3 above, the proteins are recovered from the cell culture and purified by isolating the BMP-11 proteins from other proteinaceous materials with which they are co-produced as well as from other contaminants. The purified protein may be assayed in accordance with the rat bone formation assay described in Example 6.

Purification is carried out using standard techniques known to those skilled in the art.

Protein analysis is conducted using standard techniques such as SDS-PAGE acrylamide [Laemmli, Nature 227:680 (1970)] stained with silver [Oakley, et al. Anal. Biochem. 105:361 (1980)] and by immunoblot [Towbin, et al. Proc. Natl. Acad. Sci. USA 76:4350 (1979)].

The foregoing descriptions detail presently preferred embodiments of the present invention. Numerous modifications and variations in practice thereof are expected to occur to those skilled in the art upon consideration of these descriptions. Those modifications and variations are believed to be encompassed within the claims appended hereto.

EXAMPLE 8**Tests to determine activin activity of BMP-11**

Purification is carried out using standard techniques known to those skilled in the art. It is contemplated, as with other proteins of the TGF- β superfamily, that purification may include the use of Heparin sepharose.

Protein analysis is conducted using standard techniques such as SDS-PAGE acrylamide [Laemmli, Nature 227:680 (1970)] stained with silver [Oakley, et al. Anal. Biochem. 105:361 (1980)] and by immunoblot [Towbin, et al. Proc. Natl. Acad. Sci. USA 76:4350 (1979)].

BMP-11 proteins may be further characterized by their ability to modulate the release of follicle stimulating hormone (FSH) in established in vitro bioassays using rat anterior pituitary cells as described in, for example, Vale et al,

Endocrinology, 91:562-572 (1972); Ling et al., *Nature*, 321:779-782 (1986) or Vale et al., *Nature*, 321:776-779 (1986), the disclosures of which are hereby incorporated by reference.

5 Alternatively, BMP-11 may be characterized by their ability to stimulate erythropoietin activity in the human K-562 cell line, as described by Lozzio et al., *Blood*, 45:321-334 (1975) and United States Patent No. 5,071,834, at column 15, the disclosures of which are hereby incorporated by reference.

10 In addition, BMP-11 may be characterized by their activity in cell survival assays, as described in Schubert, *Nature*, 344:868-870 (1990), the disclosure of which is incorporated by reference.

15 The foregoing descriptions, detail presently preferred embodiments of the present invention. Numerous modifications and variations in practice thereof are expected to occur to those skilled in the art upon consideration of these descriptions. Those modifications and variations are believed to be encompassed within the claims appended hereto.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: GENETICS INSTITUTE, INC.
 (B) STREET: 87 CambridgePark Drive
 (C) CITY: Cambridge
 (D) STATE: Massachusetts
 (E) COUNTRY: USA
 (F) POSTAL CODE (ZIP): 02140
 (G) TELEPHONE: 617 876-1170
 (H) TELEFAX: 617-876-5851

(ii) TITLE OF INVENTION: BMP-11 COMPOSITIONS

(iii) NUMBER OF SEQUENCES: 11

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 789 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Bos Taurus
 (B) STRAIN: Bovine Activin WC

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: 324..704

(ix) FEATURE:

(A) NAME/KEY: misc feature
 (B) LOCATION: 322..323
 (D) OTHER INFORMATION: /note= "putative 3' end of intron"

(ix) FEATURE:

(A) NAME/KEY: mat_peptide
 (B) LOCATION: 375..701

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```

AAACTGTATT TTGGGGTGAA GGTGTGAGTT AATAGATTCA CGGGACAACA AAGATGGGCT      60
GTTGTTGAGA CCTTGGGCCA AGGGGCTGAT GAGGGTCAGG TTGCCAAGAG AGAGAGAATT      120
AGGGAAGGTG AGTTTAGGGA GACATGGCTA GCTGGCAAGA AAAGTGGGTA GAAAACAGGG      180
GTTGGGGAGG GGAGCACTGG AGAAGCTCAG AAATCACTTG GTCTCTGTTC TCCTGCCCCCT      240
ACTGAGGGGC AGGTGAGAAG AAACAGGGAG TAGGAGCTCC TCGAGGCTCT ATTACATCTC      300

```

[illegible]

(2) INFORMATION FOR SEQ ID NO: 2:

(1) SEQUENCE CHARACTERISTICS:

- SEQUENCE CHARACTERISTICS:
(A) LENGTH: 126 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

His	Pro	Phe	Met	Glu	Leu	Arg	Val	Leu	Glu	Asn	Thr	Lys	Arg	Ser	Arg
-17		-15					-10					-5			
Arg	Asn	Leu	Gly	Leu	Asp	Cys	Asp	Glu	His	Ser	Ser	Glu	Ser	Arg	Cys
	1				5					10					15
Cys	Arg	Tyr	Pro	Leu	Thr	Val	Asp	Phe	Glu	Ala	Phe	Gly	Trp	Asp	Trp
				20					25					30	
Ile	Ile	Ala	Pro	Lys	Arg	Tyr	Lys	Ala	Asn	Tyr	Cys	Ser	Gly	Gln	Cys
			35					40					45		
Glu	Tyr	Met	Phe	Met	Gln	Lys	Tyr	Pro	His	Thr	His	Leu	Val	Gln	Gln
		50					55					60			
Ala	Asn	Pro	Arg	Gly	Ser	Ala	Gly	Pro	Cys	Cys	Thr	Pro	Thr	Lys	Met

65	70	75
Ser Pro Ile Asn Met Leu Tyr Phe Asn Asp Lys Gln Gln Ile Ile Tyr		
80	85	90
Gly Lys Ile Pro Gly Met Val Val Asp Arg Cys Gly Cys Ser		
100	105	

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 213 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (B) STRAIN: Human Activin WC

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 28..183

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 184..185
- (D) OTHER INFORMATION: /note= "two-thirds of codon at end of partial clone"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

TCTAGATGCT CCGGCCAGTG CGAGTAC ATG TTC ATG CAA AAA TAT CCG CAT	51
Met Phe Met Gln Lys Tyr Pro His	
1 5	
ACC CAT TTG GTG CAG CAG GCC AAT CCA AGA GGC TCT GCT GGG CCC TGT	99
Thr His Leu Val Gln Gln Ala Asn Pro Arg Gly Ser Ala Gly Pro Cys	
10 15 20	
TGT ACC CCC ACC AAG ATG TCC CCA ATC AAC ATG CTC TAC TTC AAT GAC	147
Cys Thr Pro Thr Lys Met Ser Pro Ile Asn Met Leu Tyr Phe Asn Asp	
25 30 35 40	
AAG CAG CAG ATT ATC TAC GGC AAG ATC CCT GGC ATG GTGGTGGATC	193
Lys Gln Gln Ile Ile Tyr Gly Lys Ile Pro Gly Met	
45 50	
GCTGTGGCTG CTCCGGATCC	213

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Phe Met Gln Lys Tyr Pro His Thr His Leu Val Gln Gln Ala Asn
 1 5 10 15
 Pro Arg Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro
 20 25 30
 Ile Asn Met Leu Tyr Phe Asn Asp Lys Gln Gln Ile Ile Tyr Gly Lys
 35 40 45
 Ile Pro Gly Met
 50

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: primer C to Bovine Activin WC

(ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..9
- (D) OTHER INFORMATION: /note= "Restriction site for XbaI"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

TAGTCTAGAT GCTCCGGCCA GTGCGAGTAC

30

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Primer D to Bovine Activin WC

(ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..9
- (D) OTHER INFORMATION: /note= "Restriction site for BamHI"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

TGC GGATCCG GAGCAGCCAC AGCGATCCAC

30

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: DNA inserted into pMT2 CXM

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CATGGGCAGC TCGAG

15

- (2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: DNA inserted into pMT21

- (ix) FEATURE:
(A) NAME/KEY: misc_feature
(B) LOCATION: 1..6
(D) OTHER INFORMATION: /note= "Pst restriction site"

- (ix) FEATURE:
(A) NAME/KEY: misc feature
(B) LOCATION: 15..26
(D) OTHER INFORMATION: /note= "Eco RI and XhoI restriction sites"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

CTGCAGGCGA GCCTGAATTC CTCGAGCCAT CATG

34

- (2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 68 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Portion of the EMC virus leader sequence

- (x) PUBLICATION INFORMATION:
(A) AUTHORS: Jung, S K
(C) JOURNAL: J. Virol.
(D) VOLUME: 63
(F) PAGES: 1651-1660
(G) DATE: 1989

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

CGAGGTTAAA AAACGTCTAG GCGGGGCGAA CCACGGGGAC GTGGTTTTC TTTGAAAAAC
ACGATTGC

60

68

- (2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1270 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Human BMP-11

(vii) IMMEDIATE SOURCE:

(B) CLONE: FB30.5

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..1086

(ix) FEATURE:

(A) NAME/KEY: mat_peptide

(B) LOCATION: 760..1086

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

GAG CGC TCC AGC CGG CCA GCC CCG TCC GTG GCG CCC GAG CCG GAC GGC	48
Glu Arg Ser Ser Arg Pro Ala Pro Ser Val Ala Pro Glu Pro Asp Gly	
-253 -250 -245 -240	
TGC CCC GTG TGC GTT TGG CGG CAG CAC AGC CGC GAG CTG CGC CTA GAG	96
Cys Pro Val Cys Val Trp Arg Gln His Ser Arg Glu Leu Arg Leu Glu	
-235 -230 -225	
AGC ATC AAG TCG CAG ATC TTG AGC AAA CTG CGG CTC AAG GAG GCG CCC	144
Ser Ile Lys Ser Gln Ile Leu Ser Lys Leu Arg Leu Lys Glu Ala Pro	
-220 -215 -210	
AAC ATC AGC CGC GAG GTG GTG AAG CAG CTG CTG CCC AAG GCG CCG CCG	192
Asn Ile Ser Arg Glu Val Val Lys Gln Leu Leu Pro Lys Ala Pro Pro	
-205 -200 -195 -190	
CTG CAG CAG ATC CTG GAC CTA CAC GAC TTC CAG GGC GAC GCG CTG CAG	240
Leu Gln Gln Ile Leu Asp Leu His Asp Phe Gln Gly Asp Ala Leu Gln	
-185 -180 -175	
CCC GAG GAC TTC CTG GAG GAG GAC GAG TAC CAC GCC ACC ACC GAG ACC	288
Pro Glu Asp Phe Leu Glu Glu Asp Glu Tyr His Ala Thr Thr Glu Thr	
-170 -165 -160	
GTC ATT AGC ATG GCC CAG GAG ACG GAC CCA GCA GTA CAG ACA GAT GGC	336
Val Ile Ser Met Ala Gln Glu Thr Asp Pro Ala Val Gln Thr Asp Gly	
-155 -150 -145	
AGC CCT CTC TGC TGC CAT TTT CAC TTC AGC CCC AAG GTG ATG TTC ACA	384
Ser Pro Leu Cys Cys His Phe His Phe Ser Pro Lys Val Met Phe Thr	
-140 -135 -130	
AAG GTA CTG AAG GCC CAG CTG TGG GTG TAC CTA CGG CCT GTA CCC CGC	432
Lys Val Leu Lys Ala Gln Leu Trp Val Tyr Leu Arg Pro Val Pro Arg	
-125 -120 -115 -110	
CCA GCC ACA GTC TAC CTG CAG ATC TTG CGA CTA AAA CCC CTA ACT GGG	480
Pro Ala Thr Val Tyr Leu Gln Ile Leu Arg Leu Lys Pro Leu Thr Gly	
-105 -100 -95	
GAA GGG ACC GCA GGG GGA GGG GGC GGA GGC CGG CGT CAC ATC CGT ATC	528
Glu Gly Thr Ala Gly Gly Gly Gly Gly Gly Arg Arg His Ile Arg Ile	

																-90																	-85																	-80																																																														
CGC	TCA	CTG	AAG	ATT	GAG	CTG	CAC	TCA	CGC	TCA	GGC	CAT	TGG	CAG	AGC	576																																																																																																
Arg	Ser	Leu	Lys	Ile	Glu	Leu	His	Ser	Arg	Ser	Gly	His	Trp	Gln	Ser																																																																																																	
			-75				-70				-65																																																																																																					
ATC	GAC	TTC	AAG	CAA	GTG	CTA	CAC	AGC	TGG	TTC	CGC	CAG	CCA	CAG	AGC	624																																																																																																
Ile	Asp	Phe	Lys	Gln	Val	Leu	His	Ser	Trp	Phe	Arg	Gln	Pro	Gln	Ser																																																																																																	
			-60				-55				-50																																																																																																					
AAC	TGG	GGC	ATC	GAG	ATC	AAC	GCC	TTT	GAT	CCC	AGT	GGC	ACA	GAC	CTG	672																																																																																																
Asn	Trp	Gly	Ile	Glu	Ile	Asn	Ala	Phe	Asp	Pro	Ser	Gly	Thr	Asp	Leu																																																																																																	
			-45				-40				-35				-30																																																																																																	
GCT	GTC	ACC	TCC	CTG	GGG	CCG	GGA	GCC	GAG	GGG	CTG	CAT	CCA	TTC	ATG	720																																																																																																
Ala	Val	Thr	Ser	Leu	Gly	Pro	Gly	Ala	Glu	Gly	Leu	His	Pro	Phe	Met																																																																																																	
			-25				-20				-15																																																																																																					
GAG	CTT	CGA	GTC	CTA	GAG	AAC	ACA	AAA	CGT	TCC	CGG	CGG	AAC	CTG	GGT	768																																																																																																
Glu	Leu	Arg	Val	Leu	Glu	Asn	Thr	Lys	Arg	Ser	Arg	Arg	Asn	Leu	Gly																																																																																																	
			-10				-5				1																																																																																																					
CTG	GAC	TGC	GAC	GAG	CAC	TCA	AGC	GAG	TCC	CGC	TGC	TGC	CGA	TAT	CCC	816																																																																																																
Leu	Asp	Cys	Asp	Glu	His	Ser	Ser	Glu	Ser	Arg	Cys	Cys	Arg	Tyr	Pro																																																																																																	
			5				10				15																																																																																																					
CTC	ACA	GTG	GAC	TTT	GAG	GCT	TTC	GGC	TGG	GAC	TGG	ATC	ATC	GCA	CCT	864																																																																																																
Leu	Thr	Val	Asp	Phe	Glu	Ala	Phe	Gly	Trp	Asp	Trp	Ile	Ile	Ala	Pro																																																																																																	
			20				25				30				35																																																																																																	
AAG	CGC	TAC	AAG	GCC	AAC	TAC	TGC	TCC	GGC	CAG	TGC	GAG	TAC	ATG	TTC	912																																																																																																
Lys	Arg	Tyr	Lys	Ala	Asn	Tyr	Cys	Ser	Gly	Gln	Cys	Glu	Tyr	Met	Phe																																																																																																	
			40				45				50																																																																																																					
ATG	CAA	AAA	TAT	CCG	CAT	ACC	CAT	TTG	GTG	CAG	CAG	GCC	AAT	CCA	AGA	960																																																																																																
Met	Gln	Lys	Tyr	Pro	His	Thr	His	Leu	Val	Gln	Gln	Ala	Asn	Pro	Arg																																																																																																	
			55				60				65																																																																																																					
GGC	TCT	GCT	GGG	CCC	TGT	TGT	ACC	CCC	ACC	AAG	ATG	TCC	CCA	ATC	AAC	1008																																																																																																
Gly	Ser	Ala	Gly	Pro	Cys	Cys	Thr	Pro	Thr	Lys	Met	Ser	Pro	Ile	Asn																																																																																																	
			70				75				80																																																																																																					
ATG	CTC	TAC	TTC	AAT	GAC	AAG	CAG	CAG	ATT	ATC	TAC	GGC	AAG	ATC	CCT	1056																																																																																																
Met	Leu	Tyr	Phe	Asn	Asp	Lys	Gln	Gln	Ile	Ile	Tyr	Gly	Lys	Ile	Pro																																																																																																	
			85				90				95																																																																																																					
GGC	ATG	GTG	GTG	GAT	CGC	TGT	GGC	TGC	TCT	TAAGGTGGGG						GATAGAGGAT						1106																																																																																										
Gly	Met	Val	Val	Asp	Arg	Cys	Gly	Cys	Ser																																																																																																							
			100				105																																																																																																									
GCCTCCCCCA																CAGACCCTAC																CCCAAGACCC																CTAGCCCTGC																CCCCATCCCC																CCAAGCCCTA																1166																
GAGCTCCCTC																CACTCTTCCC																GCGAACATCA																CACCGTTCCTC																CGACCAAGCC																GTGTGCAATA																1226																
CAACAGAGGG																AGGCAGGTGG																GAATTGAGGG																TGAGGGGTTT																GGGG																1270																																

(2) INFORMATION FOR SEQ ID NO: 11:

- (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 362 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Glu Arg Ser Ser Arg Pro Ala Pro Ser Val Ala Pro Glu Pro Asp Gly
 -253 -250 -245 -240
 Cys Pro Val Cys Val Trp Arg Gln His Ser Arg Glu Leu Arg Leu Glu
 -235 -230 -225
 Ser Ile Lys Ser Gln Ile Leu Ser Lys Leu Arg Leu Lys Glu Ala Pro
 -220 -215 -210
 Asn Ile Ser Arg Glu Val Val Lys Gln Leu Leu Pro Lys Ala Pro Pro
 -205 -200 -195 -190
 Leu Gln Gln Ile Leu Asp Leu His Asp Phe Gln Gly Asp Ala Leu Gln
 -185 -180 -175
 Pro Glu Asp Phe Leu Glu Glu Asp Glu Tyr His Ala Thr Thr Glu Thr
 -170 -165 -160
 Val Ile Ser Met Ala Gln Glu Thr Asp Pro Ala Val Gln Thr Asp Gly
 -155 -150 -145
 Ser Pro Leu Cys Cys His Phe His Phe Ser Pro Lys Val Met Phe Thr
 -140 -135 -130
 Lys Val Leu Lys Ala Gln Leu Trp Val Tyr Leu Arg Pro Val Pro Arg
 -125 -120 -115 -110
 Pro Ala Thr Val Tyr Leu Gln Ile Leu Arg Leu Lys Pro Leu Thr Gly
 -105 -100 -95
 Glu Gly Thr Ala Gly Gly Gly Gly Gly Gly Arg Arg His Ile Arg Ile
 -90 -85 -80
 Arg Ser Leu Lys Ile Glu Leu His Ser Arg Ser Gly His Trp Gln Ser
 -75 -70 -65
 Ile Asp Phe Lys Gln Val Leu His Ser Trp Phe Arg Gln Pro Gln Ser
 -60 -55 -50
 Asn Trp Gly Ile Glu Ile Asn Ala Phe Asp Pro Ser Gly Thr Asp Leu
 -45 -40 -35 -30
 Ala Val Thr Ser Leu Gly Pro Gly Ala Glu Gly Leu His Pro Phe Met
 -25 -20 -15
 Glu Leu Arg Val Leu Glu Asn Thr Lys Arg Ser Arg Arg Asn Leu Gly
 -10 -5 1
 Leu Asp Cys Asp Glu His Ser Ser Glu Ser Arg Cys Cys Arg Tyr Pro
 5 10 15
 Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp Trp Ile Ile Ala Pro
 20 25 30 35
 Lys Arg Tyr Lys Ala Asn Tyr Cys Ser Gly Gln Cys Glu Tyr Met Phe
 40 45 50
 Met Gln Lys Tyr Pro His Thr His Leu Val Gln Gln Ala Asn Pro Arg
 55 60 65
 Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys Met Ser Pro Ile Asn
 70 75 80

Met Leu Tyr Phe Asn Asp Lys Gln Gln Ile Ile Tyr Gly Lys Ile Pro
85 90 95

Gly Met Val Val Asp Arg Cys Gly Cys Ser
100 105

What is claimed is:

1. An isolated DNA sequence encoding a BMP-11 protein.
2. The DNA sequence of claim 1 wherein said DNA is selected from the group consisting of:
 - (a) nucleotide #375 or 390 to #704 of SEQ ID NO:1;
 - (b) nucleotides #760 or 775 to # 1086 of SEQ ID NO:10; and
 - (c) sequences which hybridize thereto under stringent hybridization conditions and encode a protein which exhibits BMP-11 activity.
3. The DNA sequence of claim 1 wherein said DNA is selected from the group consisting of:
 - (a) nucleotides encoding for amino acids 1 or 6 to 109 of SEQ ID NO:2;
 - (b) nucleotides encoding for amino acids 1 or 6 to 109 of SEQ ID NO:11; and
 - (c) sequences which hybridize thereto under stringent hybridization conditions and encode a protein which exhibits BMP-11 activity.
4. A host cell transformed with the DNA sequence of claim 1.
5. An isolated DNA molecule having a sequence encoding an BMP-11 protein which exhibits BMP-11 activity, said DNA molecule comprising the DNA sequence selected from the group consisting of:
 - (a) nucleotide # 375 to # 704 of SEQ ID NO:1;
 - (b) nucleotide # 760 to # 1086 of SEQ ID NO:10; and
 - (c) naturally occurring allelic sequences and equivalent degenerative codon sequences of (a) or (b).
6. A host cell transformed with the DNA molecule of claim 5.
7. A vector comprising a DNA molecule of claim 5 in operative association with an expression control sequence therefor.
8. A host cell transformed with the vector of claim 7.
9. An isolated DNA molecule encoding a BMP-11 protein, said DNA molecule comprising nucleotide # 375 to # 704 of SEQ ID NO:1 or nucleotides # 760 to # 1086 of SEQ ID NO:10.
10. A vector comprising a DNA molecule of claim 9 in operative association with an expression control sequence therefor.
11. A host cell transformed with the vector of claim 10.

12. A method for producing a purified BMP-11, protein said method comprising the steps of:
- (a) culturing a host cell transformed with a DNA molecule comprising the nucleotide sequence encoding a BMP-11 protein; and
 - (b) recovering and purifying said BMP-11 protein from the culture medium.
13. The method of claim 12, wherein said host cell is a transformed with a DNA molecule comprising a DNA coding sequence selected from the group consisting of:
- (a) nucleotide # 375 to # 704 of SEQ ID NO: 1;
 - (b) nucleotide # 760 to # 1086 of SEQ ID NO: 10; and
 - (c) naturally occurring allelic sequences and equivalent degenerative codon sequences of (a) and (b).
14. The method of claim 13, wherein said host cell is a mammalian cell and the DNA molecule further comprises a DNA sequence encoding a suitable propeptide 5' and linked in frame to the DNA coding sequence.
15. A purified BMP-11 polypeptide comprising the amino acid sequence from amino acid # 1 to # 109 as set forth in SEQ ID NO: 2.
16. A purified BMP-11 polypeptide comprising the amino acid sequence from amino acid # 1 to #109 as set forth in SEQ ID NO: 11.
17. A purified BMP-11 polypeptide of claim 16 wherein said polypeptide is a dimer wherein each subunit comprises at least the amino acid sequence from amino acid # 1 to # 109 of SEQ ID NO: 11.
18. A purified BMP-11 protein produced by the steps of
- (a) culturing a cell transformed with a DNA molecule comprising the nucleotide sequence from nucleotide # 375 to # 704 as shown in SEQ ID NO: 1; and
 - (b) recovering and purifying from said culture medium a protein comprising the amino acid sequence from amino acid # 1 to amino acid # 109 of SEQ ID NO: 2.
19. A purified BMP-11 protein produced by the steps of
- (a) culturing a cell transformed with a DNA molecule comprising the nucleotide sequence from nucleotide # 760 to

1086 as shown in SEQ ID NO: 10; and

(b) recovering and purifying from said culture medium a protein comprising the amino acid sequence from amino acid # 1 to amino acid # 109 as shown in SEQ ID NO: 11.

20. A purified BMP-11 polypeptide according to claim 16, wherein said polypeptide is a dimer wherein one subunit comprises at least the amino acid sequence from amino acid #1 to amino acid #109 of SEQ ID NO: 11, and one subunit comprises and amino acid sequence for a bone morphogenetic protein selected from the group consisting of BMP-1, BMP-2, BMP-3, BMP-4, BMP-5, BMP-6, BMP-7, BMP-8, BMP-9 and BMP-10.

21. A pharmaceutical composition comprising an effective amount of the BMP-11 polypeptide of claim 15 in admixture with a pharmaceutically acceptable vehicle.

22. A pharmaceutical composition comprising an effective amount of the BMP-11 polypeptide of claim 16 in admixture with a pharmaceutically acceptable vehicle.

23. A pharmaceutical composition comprising an effective amount of a BMP-11 protein in dimeric form with another inhibin- β protein, in admixture with a pharmaceutically acceptable vehicle.

24. A pharmaceutical composition comprising an effective amount of a BMP-11 protein in dimeric form with an inhibin- α protein, in admixture with a pharmaceutically acceptable vehicle.

25. A pharmaceutical composition comprising an effective amount of a BMP-11 protein in dimeric form with a BMP protein, in admixture with a pharmaceutically acceptable vehicle.

26. A pharmaceutical composition comprising an effective amount of a BMP-11 protein in dimeric form, wherein said composition further comprises at least one bone morphogenetic protein.

27. A chimeric DNA molecule comprising a DNA sequence encoding a propeptide from a member of the TGF- β superfamily of proteins linked in frame to a DNA sequence encoding an BMP-11 polypeptide.

INTERNATIONAL SEARCH REPORT

I. National Application No
PCT/US 94/05288

A. CLASSIFICATION OF SUBJECT MATTER <p style="margin-left: 40px;">C 12 N 15/12, C 12 P 21/02, C 07 K 13/00, C 12 N 5/10, A 61 K 37/02</p> <p style="margin-left: 20px;">According to International Patent Classification (IPC) or to both national classification and IPC 5</p>		
B. FIELDS SEARCHED <p style="margin-left: 20px;">Minimum documentation searched (classification system followed by classification symbols) C 12 N, C 12 P, C 07 K, A 61 K</p> <p style="margin-left: 20px;">Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched</p> <p style="margin-left: 20px;">Electronic data base consulted during the international search (name of data base and, where practical, search terms used)</p>		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO, A1, 92/09 697 (CELTRIX LABORATORIES) 11 June 1992 (11.06.92), claims.	1-27
P, A	WO, A1, 93/09 229 (GENETICS INSTITUTE) 13 May 1993 (13.05.93), claims.	1-27
A	WO, A1, 88/00 205 (GENETICS INSTITUTE) 14 January 1988 (14.01.88), pages 61-73.	1-16
A	US, A, 5 166 058 (WANG et al.) 24 November 1992 (24.11.92), claims.	1-16
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input type="checkbox"/> Patent family members are listed in annex.		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents:</p> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>* & * document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <p style="text-align: center;">01 September 1994</p>		Date of mailing of the international search report <p style="text-align: center;">13 -10- 1994</p>
Name and mailing address of the ISA <p style="margin-left: 40px;">European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016</p>		Authorized officer <p style="text-align: center;">WOLF e.h.</p>

INTERNATIONAL SEARCH REPORT

-2-

International Application No

PCT/US 94/05288

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	<p>WO, A1, 92/05 199 (GENETICS INSTITUTE) 02 April 1992 (02.04.92), claims.</p>	<p>1-16, 21,22</p>
A	<p>US, A, 5 187 076 (WOZNEY et al.) 16 February 1993 (16.02.93), claims.</p>	<p>1-16</p>
A	<p>WO, A1, 91/18 098 (GENETICS INSTITUTE) 28 November 1991 (28.11.91), claims.</p>	<p>1-16, 21,22</p>
A	<p>WO, A1, 93/00 432 (GENETICS INSTITUTE) 07 January 1993 (07.01.93), claims.</p>	<p>1-16, 21,22</p>

ANHANG

zum internationalen Recherchen-
bericht über die internationale
Patentanmeldung Nr.

ANNEX

to the International Search
Report to the International Patent
Application No.

ANNEXE

au rapport de recherche inter-
national relatif à la demande de brevet
international n°

PCT/US 94/05288 SAE 91095

In diesem Anhang sind die Mitglieder
der Patentfamilien der im obenge-
nannten internationalen Recherchenbericht
angeführten Patentedokumente angegeben.
Diese Angaben dienen nur zur Unter-
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family
members relating to the patent documents
cited in the above-mentioned inter-
national search report. The Office is
in no way liable for these particulars
which are given merely for the purpose
of information.

La présente annexe indique les
membres de la famille de brevets
relatifs aux documents de brevets cités
dans le rapport de recherche inter-
national visée ci-dessus. Les renseigne-
ments fournis sont donnés à titre indica-
tif et n'engagent pas la responsabilité
de l'Office.

Im Recherchenbericht angeführtes Patentedokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
WD A1 9209697	11-06-92	AU A1 91419/91 AU B2 651421 CA AA 2071912 EP A1 513334 EP A4 513334 JP T2 5505404	25-06-92 21-07-94 31-05-92 19-11-92 04-08-93 12-08-93
WD A1 9309229	13-05-93	AU A1 30622/92 EP A1 612348 MX A1 9206315	07-06-93 31-08-94 29-07-93
WD A1 8800205	14-01-88	AU A1 77835/87 AU B2 613314 DK A0 1062/88 DK A 1062/88 EP A1 313578 EP A4 313578 ES AF 2007625 GR A 871028 IL A0 83003 JP T2 2500241 NO A 880701 NO A0 880701 NZ A 220894 PT A 85225 PT B 85225 US A 4877864 US A 5106748 US A 5141905 US A 5187076 US A 5013649 US A 5116738 US A 5166058 US A 5108922 ZA A 8704681 AU A1 34487/89 AU B2 645244 EP A1 408649 EP A4 408649 JP T2 3503649 WD A1 8910409 AU A1 53577/90 AU B2 624940 CA AA 2030518 EP A1 429570 JP T2 3505098 WD A1 9011366	29-01-88 01-08-91 29-02-88 28-04-88 03-05-89 03-10-90 01-07-89 11-01-88 20-12-87 01-02-90 17-02-88 17-02-88 28-05-90 01-08-87 30-03-90 31-10-89 21-04-92 25-08-92 16-02-93 07-05-91 26-05-92 24-11-92 28-04-92 27-04-88 24-11-89 13-01-94 23-01-91 27-11-91 15-08-91 02-11-89 22-10-90 25-06-92 29-09-90 05-06-91 07-11-91 04-10-90
US A 5166058	24-11-92	AU A1 77835/87 AU B2 613314 DK A0 1062/88 DK A 1062/88 EP A1 313578 EP A4 313578 ES AF 2007625 GR A 871028 IL A0 83003 JP T2 2500241 NO A 880701 NO A0 880701 NZ A 220894 PT A 85225 PT B 85225 WD A1 8800205	29-01-88 01-08-91 29-02-88 28-04-88 03-05-89 03-10-90 01-07-89 11-01-88 20-12-87 01-02-90 17-02-88 17-02-88 28-05-90 01-08-87 30-03-90 14-01-88

ZA A	8704681	27-04-88
US A	5013649	07-05-91
US A	5106748	21-04-92
US A	5108922	28-04-92
US A	5116738	26-05-92
US A	5141905	25-08-92
US A	5187076	16-02-93
US A	4877864	31-10-89
AU A1	34487/89	24-11-89
AU B2	645244	13-01-94
EP A1	408649	23-01-91
EP A4	408649	27-11-91
JP T2	3503649	15-08-91
WD A1	8910409	02-11-89
AU A1	53577/90	22-10-90
AU B2	624940	25-06-92
CA AA	2030518	29-09-90
EP A1	429570	05-06-91
JP T2	3505098	07-11-91
WD A1	9011366	04-10-90

WD A1	9205199	02-04-92	EP A1	550625	14-07-93
-------	---------	----------	-------	--------	----------

US A	5187076	16-02-93	AU A1	77835/87	29-01-88
------	---------	----------	-------	----------	----------

AU B2	613314	01-08-91
DK A0	1062/88	29-02-88
DK A	1062/88	28-04-88
EP A1	313578	03-05-89
EP A4	313578	03-10-90
ES AF	2007625	01-07-89
GR A	871028	11-01-88
IL A0	83003	20-12-87
JP T2	2500241	01-02-90
NO A	880701	17-02-88
NO A0	880701	17-02-88
NZ A	220894	28-05-90
PT A	85225	01-08-87
PT B	85225	30-03-90
WD A1	8800205	14-01-88
ZA A	8704681	27-04-88
US A	5013649	07-05-91
US A	5106748	21-04-92
US A	5108922	28-04-92
US A	5116738	26-05-92
US A	5141905	25-08-92
US A	5166058	24-11-92
US A	4877864	31-10-89
AU A1	34487/89	24-11-89
AU B2	645244	13-01-94
EP A1	408649	23-01-91
EP A4	408649	27-11-91
JP T2	3503649	15-08-91
WD A1	8910409	02-11-89
AU A1	53577/90	22-10-90
AU B2	624940	25-06-92
CA AA	2030518	29-09-90
EP A1	429570	05-06-91
JP T2	3505098	07-11-91
WD A1	9011366	04-10-90

WD A1	9118098	28-11-91	CA AA	2082941	17-11-91
			EP A1	536186	14-04-93
			JP T2	6500991	27-01-94

WD A1	9300432	07-01-93	AU A1	22699/92	25-01-93
			AU B2	652472	25-08-94
			CA AA	2108770	26-12-92
			EP A1	592562	20-04-94